

Food and Drug Administration, HHS

§ 524.1195

aeruginosa], coagulase-positive staphylococci, *Enterococcus faecalis*, *Proteus mirabilis*, and beta-hemolytic streptococci).

[66 FR 712, Jan. 4, 2001, as amended at 68 FR 15370, Mar. 31, 2003]

§ 524.1140 Imidacloprid and ivermectin.

(a) *Specifications.* The product is available in unit applicator tubes containing 0.4, 1.0, 2.5, or 4.0 milliliters (mL). Each mL of solution contains 100 milligrams (mg) imidacloprid and 800 micrograms (µg) ivermectin.

(b) *Sponsor.* See No. 000859 in § 510.600(c) of this chapter.

(c) *Conditions of Use in Dogs*—(1) *Amount.* The recommended minimum dosage is 4.5 mg/pound (lb) (10 mg/kilogram (kg)) of imidacloprid and 36.4 µg/lb (80 µg/kg) of ivermectin, topically once a month.

(2) *Indications for Use.* For the prevention of heartworm disease caused by *Dirofilaria immitis*; kills adult fleas and is indicated for the treatment of flea infestations (*Ctenocephalides felis*).

(3) *Limitations.* Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[67 FR 78685, Dec. 26, 2002]

§ 524.1193 Ivermectin pour-on.

(a) *Specifications.* Each milliliter (mL) of solution contains 5 milligrams of ivermectin.

(b) *Sponsors.* See sponsors in § 510.600(c) of this chapter for use as in paragraph (e) of this section.

(1) No. 050604 for use as in paragraphs (e)(1), (e)(2)(i), (e)(2)(iii), and (e)(3) of this section.

(2) Nos. 051259, 051311, 055529, 058829, 059130, and 066916 for use as in paragraphs (e)(1), (e)(2)(i), (e)(2)(ii), and (e)(3) of this section.

(c) *Related tolerances.* See § 556.344 of this chapter.

(d) *Special considerations.* See § 500.25 of this chapter.

(e) *Conditions of use in cattle*—(1) *Amount.* One mL per 22 pounds (0.5 milligram per kilogram) of body weight applied topically to the back of the animal.

(2) *Indications for use*—(i) It is used for the treatment and control of: Gas-

trointestinal roundworms (adults and fourth-stage larvae) *Ostertagia ostertagi* (including inhibited stage), *Haemonchus placei*, *Trichostrongylus axei*, *T. colubriformis*, *Cooperia oncophora*, *C. punctata*, *C. surnabada*, *Oesophagostomum radiatum*; (adults) *Strongyloides papillosus*, *Trichuris* spp.; lungworms (adults and fourth-stage larvae) *Dictyocaulus viviparus*; cattle grubs (parasitic stages) *Hypoderma bovis*, *H. lineatum*; mites *Sarcoptes scabiei* var. *bovis*; lice *Linognathus vituli*, *Haematopinus eurysternus*, *Damalinia bovis*, *Solenopotes capillatus*; and horn flies *Haematobia irritans*.

(ii) It controls infections and prevents reinfection with *O. ostertagi*, *O. radiatum*, *H. placei*, *T. axei*, *C. punctata*, and *C. oncophora* for 14 days after treatment.

(iii) It controls infections and prevents reinfection with *O. radiatum* and *D. viviparus* for 28 days after treatment, *C. punctata* and *T. axei* for 21 days after treatment, *H. placei*, *C. oncophora*, and *C. surnabada* for 14 days after treatment, and *D. bovis* for 56 days after treatment.

(3) *Limitations.* Do not treat cattle within 48 days of slaughter. Because a withdrawal time in milk has not been established, do not use in female dairy cattle of breeding age. A withdrawal period has not been established for this product on preruminating calves. Do not use on calves to be processed for veal.

[55 FR 50551, Dec. 7, 1990, as amended at 62 FR 38908, July 21, 1997; 62 FR 63271, Nov. 28, 1997; 63 FR 44385, Aug. 19, 1998; 66 FR 13236, Mar. 5, 2001; 66 FR 63165, Dec. 5, 2001; 68 FR 3817, Jan. 27, 2003; 68 FR 4713, Jan. 30, 2003; 69 FR 501, Jan. 6, 2004; 69 FR 62181, Oct. 25, 2004]

§ 524.1195 Ivermectin otic suspension.

(a) *Specifications.* Each tube contains 0.5 milliliter (mL) of a 0.01 percent suspension of ivermectin.

(b) *Sponsor.* See No. 065274 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount.* Administer the contents of one 0.5-mL tube topically into each external ear canal.

(2) *Indications for use.* For the treatment of adult ear mite (*Otodectes cynotis*) infestations in cats and kittens 4 weeks of age and older. Effectiveness